

Phase Transitions Pt 4

Spatial correlations:

Important physical quantity in science, especially phase transitions are spatial correlations. This means if something is true at place \vec{r} , how likely is it to be true at place $\vec{r} + \vec{r}'$. This was like $g(r)$, how much more likely are we to find a particle than any a given distance away from something we also measured this as

$$\langle \rho(\vec{r}) \rho(\vec{r}') \rangle \quad (\text{or } \langle \rho(\vec{r}) \rho(\vec{r} + \vec{r}') \rangle)$$


In a lattice model, we compute the following related quantity

$$C_{ij} = \langle (s_i - \langle s_i \rangle) (s_j - \langle s_j \rangle) \rangle = \langle s_i s_j \rangle - \langle s_i \rangle \langle s_j \rangle$$

called the spin-spin correlation function

Goes to zero when $\langle s_i s_j \rangle \rightarrow \langle s_i \rangle \langle s_j \rangle$,
 which happens when uncorrelated,
 eg $|j-i| \gg 0$, sometimes normalize

$$C_{ij}, \quad \tilde{C}_{ij} = \frac{\langle s_i s_j \rangle - \langle s_i \rangle \langle s_j \rangle}{\langle s_i^2 \rangle - \langle s_i \rangle^2} \quad \text{st}$$



spins correlated w/ spin 1, eg

$$= \sum_{j=2}^{\infty} C_{ij} \quad (\text{could have chosen any spin})$$

This is connected to the susceptibility

$$\chi = \frac{1}{N} \langle (SM)^2 \rangle \quad \text{where} \quad SM = \sum_{i=1}^N (s_i - \langle s_i \rangle)$$



$$= \frac{1}{N} \sum_{i,j} \langle s_i s_j \rangle - \langle s_i \rangle \langle s_j \rangle$$

$$= \frac{N}{N} \cdot \sum_j \langle s_i s_j \rangle - \langle s_i \rangle \langle s_j \rangle = \sum_j C_{ij}$$

But we already saw that χ diverges @ a phase transition, so this is a signature of "long range correlations"

2 ways this can diverge

1) 2 phase coexistence



as these get more sharp w/ increasing N , variance gets maximized

Near critical point, no distinction between two phases, so the divergence becomes large means correlations become long range


usually $G(r) \approx C_1 r \sim \frac{e^{-r/\xi}}{r^{d-2+\nu}}$ fit to

and as critical point is approached $\xi \sim |T - T_c|^{-\nu}$

The fact that this length scale gets big means the system looks the same on small & large length scales, which leads to the renormalization group idea!

Renormalization Group

"coarse grain" over section of system
 try to have partition function look the same



then eg $H = \sum_{\langle ij \rangle} J s_i s_j \rightarrow \sum_{\langle i', j' \rangle} J' s_{i'} s_{j'}$ (zero field)

Then you can repeat this procedure & iterate

If $J \rightarrow J'$ process converges, then this is at a "fixed point" and it corresponds to a phase transition.

From the equations that generate $J \rightarrow J'$, we can get the critical exponents too!

We'll illustrate in 1d Ising model (no spontaneous mag transition) & discuss the result in 2d [1 of different possible procedures]

@ zero field

$$Q(K, N) = \sum_{s_1, \dots, s_N = \pm 1} \exp[K (s_1 s_2 + s_2 s_3 + \dots + s_N s_1)]$$

\swarrow $k = \beta J$

In mean field, averaged over all other

degrees of freedom but 1. Here,
 remove finite degrees of freedom.
 Eg: Sum over even spins

$$\overline{1} \quad \overline{2} \quad \overline{3} \quad \overline{4} \quad \overline{5} \quad \overline{6} \rightarrow \overline{1} \quad \overline{3} \quad \overline{5}$$

$$Q(k, N) = \sum_{s_1, s_3, \dots} \left(\exp[k(s_1 + s_3)] + \exp[-k(s_1 + s_3)] \right) \\
\times \left(\exp[k(s_3 + s_5)] + \exp[-k(s_3 + s_5)] \right) \\
\times \dots$$

want to express in same as original form,
 if possible, but w/ new k' (new
 inverse temperature, or β)

would be true if $e^{k(s+s')} + e^{-k(s+s')} = f(k) e^{k'ss'}$
 for all ss'

then $Q(k, N) = f(k)^{N/2} Q(k', N/2)$
 [Kadanoff transformation]

finding this is possible in 1d
[in higher d, have to approximate]

if s, s' in same direction

$$e^{zk} + e^{-zk} = f(k) e^{k'} \quad (1)$$

if opposite

$$2 = f(k) e^{-k'} \Rightarrow f(k) = 2e^{k'} \quad (2)$$

can solve for k' & $f(k)$

$$(2) \rightarrow (1) \quad 2e^{2k'} = e^{zk} + e^{-zk}$$
$$\Rightarrow k' = \frac{1}{2} \log(\cosh(2k))$$

plugging back in to 2

$$f(k) = 2 \cosh^{1/2}(2k)$$

Consider $\ln Q$ (a sort of free energy)

expect to grow $\propto N$. define $g(k) = \frac{1}{N} \ln Q$

[intensive free energy]

$$g(k) = \frac{1}{N} \cdot \left[\frac{N}{2} \ln f(k) + \ln Q(k', N/2) \right]$$

$\frac{1}{2} g(k')$

⇒ recursion:

$$\begin{aligned}g(k') &= 2g(k) - \ln f(k) \\ &= 2g(k) - \ln(2 \sqrt{\cosh(2k)})\end{aligned}$$

If we know $Q(k)$ for 1 value of k , we can find it for other values

In this renormalization, k' is always less than k
[e.g. $\frac{1}{2} \ln \cosh(2k) = k' < k$]

Alternatively, swap k & k'

$$k = \frac{1}{2} \cosh^{-1}(\exp(2k'))$$

& solve for $g(k)$

$$\begin{aligned}g(k) &= \frac{1}{2}g(k') + \frac{1}{2} \ln 2 + \frac{1}{2} \ln(\cosh(\cosh^{-1}(\exp(2k')))) \\ &= \frac{1}{2}g(k') + \frac{1}{2} \ln 2 + k'/2\end{aligned}$$

What do we do with this? For small k' , e.g.

$k' = 0.01$, high temp or low J means spins

close to uncorrelated $\mathcal{Q}(k', N) \approx 2^N$, $g(k') = \ln 2$

$$\Rightarrow \begin{cases} k = 0.100534 \\ g(k) = 0.698147 \end{cases}$$

$$\approx 0.693$$

keep iterating & k keeps growing
 during the iteration, $g(k)$ gets closer
 & closer to the exact $g(k)$ for that
 value of k [in 1d], can compute exactly

For no field & large N , need seen

$$Q(\beta, N) = (2 \cosh(\beta J))^N$$


$$= (e^k + e^{-k})^N$$

$$\ln Q/N = \ln(e^k + e^{-k}) = k + \ln(1 + e^{-2k})$$

for large k , $g(k) \approx k$

can start w/ e.g. $g(10) \approx 10$ &
 iterate other equations to get smaller k
 results

Errors apparently grow in this procedure,
 but the picture is



for $k=0$ or $k=\infty$, params don't change "fixed point"

In 2d

$x \leftarrow \leftarrow \leftarrow \leftarrow x \rightarrow \rightarrow \rightarrow \rightarrow x$
 $k=0 \qquad \qquad \qquad k_c \qquad \qquad \qquad k=\infty$

Unstable fixed point at

$k_c = 0.50698$ (for a similar problem
See Chandler Ch 5)

where exact value $J/k_B T_c = k_c = 0.44089$

See book pg 261 that shows # iterations needed as get closer & closer to k_c grows, connecting to a growing length scale as T_c approached